

Symmetries of the Weyl tensor in Bianchi V spacetimes

A. R. Kashif^{1*}, K. Saifullah^{2†} and G. Shabbir³

¹*Department of Mathematical Sciences, University of Aberdeen, Aberdeen AB24 3UE, Scotland, UK*

²*School of Mathematical Sciences, Queen Mary, University of London, London, UK*

³*Faculty of Engineering Sciences, GIK Institute of Engineering Sciences and Technology, Topi, Swabi, NWFP, Pakistan*
Electronic address: kashmology@yahoo.com, saifullah@qau.edu.pk, shabbir@giki.edu.pk

ABSTRACT: Symmetries of geometrical and physical quantities in general relativity provide important information about the curvature structure of the spacetimes. Symmetries of the curvature and the Weyl tensors, known as curvature and Weyl collineations respectively, are two of such important symmetries. Some results on these symmetries for Bianchi type V spacetimes are discussed.

* *On leave from:* College of EME, National University of Sciences and Technology, Rawalpindi, Pakistan.

† *On leave from:* Centre for Advanced Mathematics and Physics, National University of Sciences and Technology, Rawalpindi, Pakistan, *and* Department of Mathematics, Quaid-i-Azam University, Islamabad, Pakistan.

Symmetries of tensors in general relativity – Killing vectors and collineations – play an important role in understanding not only the geometric structure of the underlying spaces but their physical properties as well. They have been used in finding new solutions of Einstein’s Field Equations (EFEs), classifying these solutions and by virtue of Noether’s theorem constructing the conservation laws for the spacetime. Thus, invariance under the Lie transport of the metric, Ricci and energy momentum tensors define Killing vectors (KVs), Ricci collineations (RCs) and matter collineations (MCs), respectively [1, 2]. Curvature collineations (CCs) which are symmetries of the Riemann tensor are significant for studying the curvature structure of spacetimes [3, 2]. The Weyl tensor, $\mathbf{C}$, is fundamental in understanding the purely gravitational field for a spacetime with the matter content removed [4]. Its local symmetries, Weyl collineations (WCs) [3, 5, 6, 7], are of particular interest since it is conformally invariant [8]. Mathematically, WCs are given by

$$\mathcal{L}_{\mathbf{X}} \mathbf{C} = 0, \quad (1)$$

where $\mathcal{L}_{\mathbf{X}}$ is the Lie derivative along the vector field $\mathbf{X}$. In component form this becomes

$$C_{bcd,f}^a X^f + C_{fcd}^a X_{,b}^f + C_{bfd}^a X_{,c}^f + C_{bcf}^a X_{,d}^f - C_{bcd}^f X_{,f}^a = 0,$$

where comma denotes the partial derivative. This is a system of 20 nonlinear partial differential equations as compared to the collineations of rank two tensors (RCs and MCs, for example) which are systems of 10 equations.

On account of its algebraic symmetries we can write the 4th rank Weyl tensor (and the curvature tensor) of 4-dimensions in the form of a 6 dimensional matrix, whose rank gives the rank of the tensor. Further, while the metric tensor cannot be degenerate, the other tensors can be and hence give way to the possibility of infinite degrees of freedom (i.e. infinite dimensional Lie algebras) as well. The KVs of a space form a subset of all other collineations but the inclusion relationship between the symmetries of two fourth rank tensors, CCs and WCs, when both are finite is yet to be established. While it is known [5] that CCs can be properly contained in WCs when both are finite, no spacetime is known to the present authors which admits CCs which are not WCs and yet both are finite. On the other hand, there is no proof available that this is not possible. The Schwarzschild interior spacetime, for example, is Petrov type O [1] and thus every vector field is a WC while CCs are finite. The Reissner-Nordstrom spacetime is of Petrov type D and both the WCs and CCs are finite and equal. But when we take pressure as constant in the Schwarzschild interior we see that the WCs are properly contained in infinitely many CCs. For vacuum spacetimes with zero cosmological term, however, the Ricci tensor, $\mathbf{R}$, is zero and WCs and CCs coincide because the Weyl tensor reduces to the curvature tensor.

Enumeration of all Lie groups is useful in mathematics as well as in physics. The G_3 , for example, were originally enumerated by Bianchi which were divided into nine types, *Bianchi I* to *Bianchi IX* [1].

Let us consider the Bianchi type V spacetimes which admit three KVs given by [1]

$$\begin{aligned} K^1 &= \frac{\partial}{\partial y} , \\ K^2 &= \frac{\partial}{\partial z} , \\ K^3 &= \frac{\partial}{\partial x} - y \frac{\partial}{\partial y} - z \frac{\partial}{\partial z} . \end{aligned}$$

These spacetimes in $(t, x, y, z,)$ coordinates can be written as

$$ds^2 = -dt^2 + A(t)^2 dx^2 + B(t)^2 dy^2 + (C(t)^2 + x^2 B(t)^2) dz^2 - 2xB(t)^2 dydz. \quad (2)$$

Following a well known procedure [9, 10] the components C_{abcd} of the Weyl tensor for these spacetimes can be written as the 6×6 matrix

$$C_{abcd} = \begin{pmatrix} C_{1010} & 0 & 0 & 0 & 0 & C_{1023} \\ 0 & C_{2020} & C_{2030} & 0 & C_{1320} & 0 \\ 0 & C_{2030} & C_{3030} & C_{1230} & C_{1330} & 0 \\ 0 & 0 & C_{1230} & C_{1212} & C_{1213} & 0 \\ 0 & C_{1320} & C_{1330} & C_{1213} & C_{1313} & 0 \\ C_{1023} & 0 & 0 & 0 & 0 & C_{2323} \end{pmatrix} .$$

Similarly the curvature tensor can also be written as a 6×6 matrix. If its rank is greater than or equal to 4 then the Lie algebra of CCs is finite dimensional [11]. Now, the rank of the Weyl matrix is always even and if it is 6 or 4 the Weyl symmetry trivially reduces to the conformal symmetry [9]. Further, we note that [6] it cannot have rank 2 . Thus we conclude that these spacetimes do not admit non-trivial WCs.

Acknowledgments

ARK and KS acknowledge a research grant from the Higher Education Commission of Pakistan. They are also thankful to the National University of Sciences and Technology, Pakistan for travel support to participate in MG11, Berlin, 2006.

References

- [1] H. Stephani, D. Kramer, M. A. H. MacCallum, C. Hoenselaers and E. Herlt, *Exact Solutions of Einstein's Field Equations* (Cambridge University Press, 2003).

- [2] G. S. Hall, *Symmetries and Curvature Structure in General Relativity* (World Scientific, 2004).
- [3] G. H. Katzin, J. Levine and W. R. Davis, *J. Math. Phys.* **10**, 617 (1969).
- [4] R. Penrose and W. Rindler, *Spinors and Spacetime* (Cambridge University Press, 1986).
- [5] I. Hussain, A. Qadir and K. Saifullah, *Int. J. Mod. Phys. D* **14**, 1431 (2005).
- [6] G. Shabbir and A. R. Kashif, *A note on proper Weyl collineations in Bianchi V spacetimes*, (Submitted for publication).
- [7] G. S. Hall, *Gen. Rel. Grav.* **32**, 933 (2000).
- [8] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Spacetime* (Cambridge University Press, 1973).
- [9] G. S. Hall, *Gravitation & Cosmology* **2**, 270 (1996).
- [10] G. Shabbir, *Class. Quantum Grav.* **21**, 339 (2004).
- [11] A. H. Bokhari, A. R. Kashif and A. Qadir, *Gen. Rel. Grav.* **35**, 1059 (2003); A. R. Kashif, Ph.D. Thesis, Quaid-i-Azam University, Islamabad (2003).